

MENTAL HEALTH AND SUICIDE PREVENTION

LUNDBECK'S RECOMMENDATIONS AND COMMITMENTS



If you are thinking of suicide or are in immediate danger, please contact your local emergency services, your doctor and/or your nearest mental health crisis center. You can find a list of crisis centres around the world here:
www.iasp.info/resources/Crisis_Centres/



CONTENT

3

KEY MESSAGES AND GLOSSARY

4

SUICIDE PREVENTION IS A GLOBAL IMPERATIVE

5

SUICIDAL BEHAVIOUR IS COMPLEX YET IT IS PREVENTABLE

Suicide rate per 100,000 population by WHO region, 2016

Suicide socio-demographics and people at risk

Suicide risk factors

Suicide protective factors

Age-standardized suicide rates (per 100,000 population), both sexes, 2016

Suicide Prevention: DOs and DON'Ts

Suicide warning signs

8

10 RECOMMENDATIONS FOR A MULTI-SECTORAL APPROACH TO SUICIDE PREVENTION

What to say and what not to say

9

LUNDBECK'S COMMITMENT TO SUPPORTING MENTAL HEALTH PROMOTION AND SUICIDE PREVENTION STRATEGIES

KEY MESSAGES

1.


Due to high suicide rates, suicide prevention is a global imperative, for which national governments will be expected to deliver and report to the UN by 2030
2.

Although suicide rates are high and suicidal behaviour is complex, it is preventable by addressing risk factors, leveraging protective factors and improving healthcare systems
3.

Suicide is not only a health issue: it is a societal one. A multi-sectoral societal approach to national prevention plans is needed to help prevent suicides
4.

As a leader in restoring brain health, Lundbeck is committed to supporting mental health promotion and suicide prevention strategies

The presence of a mental health condition is a key risk factor: more than 90% of persons who die by suicide are associated with mental disorders¹², for example as:



The lifetime risk of suicide is estimated to be 4%¹³ in patients with mood disorders, 8% in people with alcohol dependence¹⁸, 8% in people with bipolar disorder¹⁹, and 5% in people with schizophrenia²⁰

GLOSSARY: DEFINITIONS

SUICIDAL BEHAVIOUR Range of behaviours that include suicide ideation (thinking about suicide, planning for suicide), attempting suicide and suicide itself¹

SUICIDAL IDEATION Thinking about, considering or planning suicide⁴⁹. DSM-5 includes suicidal ideation as a symptom of major depressive episodes⁵⁰


SUICIDE PLANNED ATTEMPT Not-fatal, self-directed, potentially injurious behaviour with intent to die (might not result in injury)⁵¹. DSM-5 includes suicide attempts as a symptom of major depressive episodes⁵⁰

SUICIDE The act of deliberately killing oneself¹




1.

Due to high suicide rates, suicide prevention is a global imperative, for which national governments will be expected to deliver and report to the UN by 2030



2.

Although suicide rates are high and suicidal behaviour is complex, it is preventable by addressing risk factors, leveraging protective factors and improving healthcare systems



SUICIDE RISK FACTORS INCLUDE

- Stigma leading to unwillingness to seek help⁷
- Difficulties in accessing treatment⁷, feelings of hopelessness²² or isolation⁷
- Loss (relational, social,work, or financial)⁷
- Previous suicide attempt(s)⁷
- The presence of a mental health condition¹²
- Chronic pain and disease²³ (cancer²⁴, diabetes²⁵, HIV/AIDS¹, Parkinson’s disease²⁶, Alzheimer’s disease²⁷)
- Child maltreatment⁷
- Family history of suicide²⁸

Suicide is preventable³


An early intervention service may be associated with reductions in the suicide rate among patients with schizophrenia-spectrum disorders during their most vulnerable period, and the benefits may persist in the long-term³⁸. Yet suicide numbers are still too high¹⁵, and likely to be underreported due to stigma, criminalization and poor surveillance systems⁵



SUICIDE PROTECTIVE FACTORS INCLUDE⁷

- Effective clinical screening and diagnosis and care for mental, physical, and substance abuse disorders
- Easy access to a variety of clinical interventions (including behavioural therapy and/or pharmacological treatment)
- Support from ongoing medical and mental health care relationships to support follow-up after discharge and treatment adherence¹
- Family and community support (connectedness)
- Cultural and religious beliefs (pending cultural and contextual practices and interpretations)
- Skills in problem solving, conflict resolution and disputes

AGE-STANDARDIZED SUICIDE RATES (PER 100,000 POPULATION), BOTH SEXES, 2016



Source: World Health Organization, 2018. Suicide rates.

SUICIDE PREVENTION: DOS¹ AND DON'Ts

DOS

Educate (yourself and others) about suicide prevention and resources while debunking myths.

When communicating, always mention where to seek help from services available 24/7.

When communicating, be mindful about celebrity suicides (focus on their life);

Consider including narratives of people who managed to cope with suicidality to inspire others.

As a primary health care provider (PHCP), be attentive to warning signs, aware of interview techniques and refer to the appropriate healthcare service/specialist.

As a healthcare professional (HCP), convey hope when diagnosing and managing a chronic or physical illness.

As a psychiatrist, ensure you are attentive to warning signs, establish a frank discussion with your patients (ask questions about suicide behaviour), follow-up on treatment adherence and refer to local peer-to-peer support groups.

As a family member, a friend, or a colleague be attentive to warning signs and encourage them to contact medical and professional support.

As a family member or a friend, establish a safe space to have discussions on how they feel and if they are thinking about suicide. Reassure them they are not alone. Remove methods of suicide and have a list of emergency contacts at hand.

DON'Ts

Fear suicide contagion and avoid talking about it.

When communicating, don't use information detailing or visualizing the method used or the location.

When communicating, don't use sensationalist language glamorizing suicide.

As an HCP, don't overlook warning signs as many of those who die by suicide have had contact with PHCPs within the month prior to the suicide.

As a HCP, avoid a tone of voice with a sense of doomed when diagnosing and managing a chronic or physical illness.

As a psychiatrist, don't fear planting a "suicide seed" in your patient's mind.

As a family member, friend or colleague don't ignore warning signs.

As a family member or a friend, don't stigmatize suicidal behavior and underestimate your role.

An important challenge on suicide prevention relates to the quality of the data collected and the risk of under-reporting (e.g. potentially due to prevailing social or religious attitudes). In some places, it is believed that

suicide is underreported by a percentage between 20% and 100%³⁹



Another big challenge is the failure of healthcare systems


to cater for people with suicidal thoughts and behaviours: GPs have increasingly limited time with each patient which can present challenges in identifying suicidal warning signs in their patients⁴⁰. When at-risk patients are identified, healthcare professionals need to exercise clinical judgement to determine the proper course of action. In the case of involuntary hospitalisation, the overall lack of hospital beds within acute psychiatry⁴¹ and fact that psychiatric hospitalisation itself presents many challenges to both provider and patient can complicate recovery. For many patients, the loss of independence, internalised and externalised stigma, and increased stress prompted by psychiatric hospitalisation must be balanced along with the need for intensive treatment services⁴²

3.

Suicide is not only a health issue: it is a societal one.
A multi-sectoral societal approach to national prevention plans is needed to help prevent suicides


LUNDBECK'S 10 RECOMMENDATIONS

POLICY




1. Ensure a national suicide prevention plan is in place and is adequately funded and monitored¹
2. Invest in national data monitoring systems and in suicidology research, e.g. on protective factors
3. Provide access to early intervention services in mental health, individualized care and treatments (including psychosocial and pharmacological interventions) as recommended by the WHO¹ and the International Association for Suicide Prevention⁴³

HEALTHCARE




4. Encourage the enrolment of medical students in the specialization of psychiatry, which is declining due to stigma of the profession, on the type of patients and of available treatments⁴⁴
5. Train (primary) healthcare professionals, to recognize, refer and manage mental and substance use disorders¹; to identify suicidal behaviour; and to convey hope to their patients with chronic disease and chronic pain⁴⁵. Ensure secondary healthcare professionals, including psychiatrists, are aware of evidence-based interventions for suicidal behaviour⁴⁶

COMMUNITY



6. Train first responders, welfare workers, educators, religious leaders¹, nursing home staff, families of people at-risk, on suicide risk factors, warning signs, adequate language and referrals to specialized care
7. Include mental health, suicide prevention and conflict resolution in school curricula
8. Put in place national media guidelines on how to report on suicide, which abide by the WHO standards and train journalists and online influencers accordingly⁴⁷
9. Reduce access to methods and secure surveillance to hot spots (e.g. bridges, rail tracks)⁴⁸
10. Support the advocacy community to drive (a) peer-to-peer support groups for attempt survivors and for families to provide a sense of connectedness; (b) suicide prevention campaigns on World Suicide Prevention Day (10 September) and World Mental Health Day (10 October) and Movember (November); (c) 24/7 helpline¹

WHAT TO SAY AND WHAT NOT TO SAY⁴⁷



DON'T SAY...	... SAY INSTEAD
Failed/unsuccessful attempt	Previous attempt OR non-fatal suicidal behaviour
Committed suicide (implies illegality, e.g. commit a crime); Completed suicide (implies accomplishment)	Died by suicide OR took his/her life
A person who failed a suicide attempt	A suicide attempt survivor

According to the WHO, despite being a preventable leading cause of death worldwide, **suicide prevention has not received the financial or human investment it needs¹**

4.

As a leader in restoring brain health, Lundbeck is committed to supporting mental health promotion and suicide prevention strategies

AT LUNDBECK, WE BELIEVE IN A MULTI-SECTORAL APPROACH TO SUICIDE PREVENTION

PATIENTS

So every person can be their best, we invest in patient education programmes globally and locally and we invest in the research, the development and patient access to treatments for depression, schizophrenia and bipolar disease.

HEALTHCARE PROFESSIONALS

We provide medical education and training on mental health promotion and suicide prevention via the Lundbeck Institute seminars, publications and online campus as well as through our disease education online platform Progress in Mind Resource Center.

FAMILY

We sponsor education programs, awareness campaigns and tools targeted at families of people with psychiatric disorders. These include information about suicide prevention.

COMMUNITY


We believe in establishing strong partnerships with the advocacy community to raise awareness and educate the media, policy-makers, healthcare professionals and the general public about mental health promotion and suicide prevention. Beyond our global partnerships, we have partnerships in the five corners of the world: from China, to the US; from Spain to Indonesia; from South Africa to Ireland.

AS AN EMPLOYER OF 5,000 PEOPLE WORLDWIDE

Lundbeck encourages every employee to become an Ambassador of change and take part of awareness raising campaigns, such as World Mental Health Day. In our affiliates, "mental health first aid" training courses (of which suicide prevention is part of) have been delivered in the UK and the US. In South Korea, our affiliate has been the first company in the country to train all its workforce as suicide prevention gatekeepers. Lundbeck Brazil partnered with the Brazilian Psychiatry Association to educate its employees on suicide prevention during "Yellow September" suicide awareness month. Employees in the US have access to the Employee Assistance Program (EAP) which provides access and referrals to mental health and support services. Our employees based in Denmark (circa 35% of Lundbeck's workforce) can take advantage of the following preventive and early care services: stress prevention courses, stress-coach scheme and psychologic help. Continuously, we will focus on the importance of early care and further strengthen the dialogue on well-being and health resilience.

Suicide is preventable³

Connectedness and a multi-sectoral approach are key to reduce suicide rates⁵. As a member of the mental health community and, considering the links between mental illness and suicidal behaviour, Lundbeck has a responsibility to people with mental disorders by providing medicines that alleviate mental disorders and to support suicide prevention policy strategies.



REFERENCES

- WHO (2014). Preventing suicide: a global imperative. World Health Organization.
- Turecki, G., & Brent, D. A. (2016). Suicide and suicidal behaviour. The Lancet, 387(10024), 1227–1239
- WHO (2014). Suicide: facts and figures World Health Organization..
- Special Advisory Committee on the Epidemic of Opioid Overdoses (2018). National report: Apparent opioid-related deaths in Canada (January 2016 to March 2018) Ottawa: Public Health Agency of Canada
- WHO (2018). Preventing suicide A community engagement toolkit.
- Holmstrand C, et al. Long-term suicide risk in no, one or more mental disorders: The Lundby Study 1947-1997. Acta Psychiatr Scand 2015;132(6):459-469.
- Centers for Disease Control and Prevention, Diseases and Conditions, Suicide (Page last reviewed: June 2018; Accessed in Jan 2019)
- J Affect Disord. 2013;151(3):821-30.; 2: Grupo de Trabajo de la Guía de Práctica Clínica de Prevención y Tratamiento de la Conducta Suicida. Guía de Práctica Clínica de Prevención y Tratamiento de la Conducta Suicida. Santiago de Compostela: Agencia de Evaluación de Tecnologías Sanitarias de Galicia (avalia-t); 2012. Guías de Práctica Clínica en el SNS: avalia-t N° 2010/02
- Donald S. Shepard et al. (2015) Suicide and Suicidal Attempts in the United States: Costs and Policy Implications. Suicide and Life-Threatening Behavior published by Wiley Periodicals.
- Kinchin I & Doran C.M (2017). The Economic Cost of Suicide and Non-Fatal Suicide Behavior in the Australian Workforce and the Potential Impact of a Workplace Suicide Prevention Strategy. Int J Environ Res Public Health v.14(4).
- The 17 Sustainable Development Goals were adopted by the United Nations Member States in 2015 to be reached by 2030. Goal #3 is to “ensure healthy lives and promote well-being for all at all ages”. The target is to reduce by one third premature mortality from non-communicable diseases through prevention and treatment and promote mental health and well-being.
- Holmstrand C., Bogren M., Mattisson C., Brådvik L. (2015). Long-term suicide risk in no, one or more mental disorders: the Lundby Study 1947–1997. Acta Psychiatr Scand. Dec; 132(6): 459–469.
- Nordentoft M, Mortensen PB, Pedersen CB. Absolute risk of suicide after first hospital contact in mental disorder. Arch Gen Psychiatry. 2011 Oct; 68(10):1058-64.
- B.A., P., V.S., P., & J.M., B. (2005). The lifetime risk of suicide in schizophrenia: a reexamination. Archives of General Psychiatry, 62(3), 247–253. 7
- Centers for Disease Control and Prevention, Diseases and Conditions, Suicide rising across the US (Page last reviewed: June 2018; Accessed in Jan 2019).
- Goodwin FK, Jamison KR (1990). Manic-Depressive Illness. New York: Oxford University Press.
- Krysinska K, Lester D. (2010). Post-traumatic stress disorder and suicide risk: a systematic review. Arch Suicide Res. 2010;14(1):1-23. doi: 10.1080/13811110903478997. Review.
- Schneider B. (2009). Substance use disorders and risk for completed suicide. Arch Suicide Res.;13(4):303–16. 1
- KL, Hawton, L, Sutton, C, Haw, J, Sinclair, L, Harriss (2005). Suicide and attempted suicide in bipolar disorder: a systematic review of risk factors. J Clin Psychiatry.; 66(6):693–704; Nordentoft M, Mortensen PB, Pedersen CB (2011). Absolute risk of suicide after first hospital contact in mental disorder. Arch Gen Psychiatry. 68(10):1058–64..
- B.A., P., V.S., P., & J.M., B. (2005). The lifetime risk of suicide in schizophrenia: a reexamination. Archives of General Psychiatry, 62(3), 247–253.
- Medical Dictionary. Suicide (accessed 27 Feb 2019)
- AT Beck, RA Steer, JS Beck, CF Newman (1993) . Hopelessness, depression, suicidal ideation, and clinical diagnosis of depression. Suicide Life Threat Behav. 1993;23:139–145
- Scott KM et al. (2010). Chronic physical conditions and their association with first onset of suicidal behavior in the world mental health surveys. Psychosom Med.
- S.Misono (2008). Incidence of Suicide in Persons With Cancer J Clin Oncol. 2008 Oct 10; 26(29): 4731–4738.
- S.Sarkar, Y.P.Singh Balhara (2014). Diabetes mellitus and suicide. Indian J Endocrinol Metab. 2014 Jul-Aug; 18(4): 468–474.
- S. Nazem, A.D. Siderowf, J.E. Duda, G.K. Brown, T.T. Have, M.B. Stern, D.Weintraub, (2008). Suicidal and Death Ideation in Parkinson’s Disease Mov Disord. 2008 Aug 15; 23(11): 1573–1579.
- Barak, D. Aizenberg D. (2002). Suicide amongst Alzheimer’s disease patients: a 10-year survey. Dement Geriatr Cogn Disord. 2002;14(2):101-3. 7
- Qin P1, Agerbo E, Mortensen PB (2002).Suicide risk in relation to family history of completed suicide and psychiatric disorders: a nested case-control study based on longitudinal registers. Lancet. 2002 Oct 12;360(9340):1126-30.
- Schmidtke et al., 2004
- Kutcher & Chehil, 2007
- Fazel S, Ramesh T, Hawton K. Suicide in prisons: an international study of prevalence and contributory factors. Lancet Psychiatry 2017 Dec;4(12):946-952.
- Centers for Disease Control (US). Suicide prevention. (accessed April 2019).
- Bilsen J. Suicide and youth: Risk factors. Front Psychiatry 2018;9:540.
- L.Vijayakumar (2015). Suicide in women. Indian J Psychiatry. 2015 Jul; 57(Suppl 2): S233- S238.
- J.B. Call, BSI, K.Shafer, Gendered Manifestations of Depression and Help Seeking Among Men, American Journal of Mens Health. 2018 Jan; 12(1): 41–51.
- Razum & Zeeb, 2004; Löhr et al., 2006; Sayil, 2006; Wohnr et al., 2006; Bursztein et al., 2009, 2010
- US National Alliance on Mental Health website, Find support, LGBTQ (accessed Feb 2019)
- Chan, S. K., Chan, S. W. Y., Pang, H. H., Yan, K. K., Hui, C. L. M., Chang, W. C., ... Chen, E. Y. H. (2018). Association of an early intervention service for psychosis with suicide rate among patients with first-episode schizophrenia-spectrum disorders. JAMA Psychiatry, 75(5), 458–464.
- J.M. Bertolote, A.Fleischmann (2020) Suicide and psychiatric diagnosis: a worldwide perspective, World of Psychiatry
- Vergier P, et al. Determinants of early identification of suicidal ideation in patients treated with antidepressants or anxiolytics in general practice: a multilevel analysis. J Affect Disord. 2007 Apr;99(1-3):253-7
- Allison S, et al. When should governments increase the supply of psychiatric beds? Mol Psychiatry. 2018 Apr;23(4):796-800. doi: 10.1038/mp.2017.139. Epub 2017 Jul 11.
- American Psychiatric Association Work Group on Suicidal Behaviors (2010). Practice Guideline for the Assessment and Treatment of Patients With Suicidal Behaviors. Psychiatry online.
- IASP Guidelines for Suicide Prevention. International Association for Suicide Prevention.
- T.Deb, G.A. Lomax, (2014). Why don’t more doctors choose a career in psychiatry?. British Medical Journal.
- W.Rutz, L. v.Knorrning, J. Wallinder (1989). Frequency of suicide on Gotland after systematic postgraduate education of general practitioners. Acta Psychiatr Scand. 1989;80:151–154
- Nordentoft M. (2011). Absolute risk of suicide after first hospital contact in mental disorder. Archives of General Psychiatry; 68: 1058-1064.
- WHO and IASP (2017). Preventing suicide: a resource for media, World Health Organization.
- Lester D, 1998. Preventing suicide by restricting access to methods for suicide. Arch Suicide Res. 1998;4:7–24
- Centers for Disease Control and Prevention (CDC), 2015
- DSM-5; Oquendo et al, 2014
- National Institute of Mental Health Information Resource Center (Last Updated: May 2018; Accessed Feb 2019)
- American Foundation for Suicide Prevention, About Suicide, Risk Factors and Warning Signs (accessed February 2019)



If you are thinking of suicide or are in immediate danger, please contact your local emergency services, your doctor and/or your nearest mental health crisis center

You can find a list of crisis centres around the world here:

www.iasp.info/resources/Crisis_Centres/



LUNDBECK
Ottiliavej 9, 2500 Valby, Denmark
+45 3630 1311 | information@lundbeck.com
www.lundbeck.com

